



Manicaland HIV STD Prevention Project Epidemic Status Report 2010

Biomedical Research & Training Institute, 4th Floor, Nicos Diamond House, Samora Machel Avenue, P.A. Box CY 1753, Harare, Zimbabwe Tel: 04735000/02 & Stand 188, Mutasa D.C., P.O. Box Hauna 3449, Zimbabwe, Tel: 0228 2230.

For

ALL SITES COVERED BY THE PROJECT

INTRODUCTION

The baseline survey for the Manicaland HIV/STD Prevention Project was carried out from July 1998 to January 2000. The first follow up survey was carried out from July 2001 to January 2003, the second follow up survey was carried out from August 2003 to August 2005, whilst the third follow up survey was carried out from August 2006 to November 2008. This sheet compares the main results on local patterns of HIV infection, experience of other sexually transmitted diseases (STDs), knowledge and awareness of HIV and other STDs, and sexual and health seeking behaviour between the baseline and the follow up surveys. Between the first and second follow-up surveys extra questions were added regarding HIV testing and treatment. See overleaf for more details.

AREAS COVERED

The survey in this area covered the following areas Katiyo, Eastern Highlands, Honde, Selbourne, Nyanga, Watsomba, Nyazura, Nyabadza/Nyahukwe, St. Theresa, Sheba, St. Killians and Bonda. In the baseline, 9,479 adults participated in the survey. In the first follow-up, 7,369 adults participated, in the second follow-up, 14,208 adults participated and, in the third follow-up, 10,525 adults participated. These numbers fluctuated due to changes in migration and changes in survey eligibility criteria.

KEY RESULTS

- 14% of men and 19% of women are currently infected with HIV.
- HIV prevalence for men and women has declined steadily since the baseline survey. However prevalence is still at a high level.
- Most people know that STDs increase the chance of HIV transmission. However, the proportion of people seeking treatment at hospitals has decreased in the last follow-up survey.
- Both men and women have reported an increase in consistent condom use with casual partners since the baseline survey.
- The proportion of people thinking that all babies born to HIV+ mothers are infected has fallen but is still more than one third.
- Fewer men but more women reported one or more new partner(s) in the last year in the most recent round of the survey.
- Stigma associated with HIV/AIDS is decreasing and most people are willing to look after a relative who has AIDS.
- The majority of people are satisfied with STD services at hospitals and clinics.
- There has been a big rise in the proportion of females having an HIV test and a smaller increase for males
- Most people are unaware that treatment for HIV/AIDS exists.